


### ***Amendments to the Claims***

The listing of claims will replace all prior versions, and listings of claims in the application.

1. (Currently amended) A composition comprising a synergistically effective combination of compounds of the formula (I)



in which

A<sup>1</sup> and A<sup>2</sup> independently of one another represent oxygen or sulfur,

X<sup>1</sup> represents N or CR<sup>10</sup>,

R<sup>1</sup> represents hydrogen or represents C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, each of which is optionally mono- or polysubstituted, where the substituents independently of one another may be selected from the group consisting of R<sup>6</sup>, halogen, cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>2</sub>-C<sub>4</sub>-alkoxycarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, (C<sub>1</sub>-C<sub>4</sub>-alkyl)-C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino and R<sup>11</sup>,

R<sup>2</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl or C<sub>2</sub>-C<sub>6</sub>-alkylcarbonyl,

R<sup>3</sup> represents hydrogen, R<sup>11</sup> or represents C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, each of which is optionally mono- or polysubstituted, where the substituents independently of one another may be selected from the group consisting of R<sup>6</sup>, halogen, cyano, nitro,

hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl, R<sup>11</sup>, phenyl, phenoxy and a 5- or 6-membered heteroaromatic ring, where each phenyl, phenoxy and 5- or 6-membered heteroaromatic ring may optionally be substituted and where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R<sup>12</sup>, or

R<sup>2</sup> and R<sup>3</sup> may be attached to one another and form the ring M,

R<sup>4</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkynyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, halogen, cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl or represents phenyl, benzyl or phenoxy, each of which is optionally mono- or polysubstituted, where the substituents independently of one another may be selected from the group consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>2</sub>-C<sub>4</sub>-haloalkenyl, C<sub>2</sub>-C<sub>4</sub>-haloalkynyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, halogen, cyano, nitro, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, C<sub>3</sub>-C<sub>6</sub>-(alkyl)cycloalkylamino, C<sub>2</sub>-C<sub>4</sub>-alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkylaminocarbonyl, C<sub>3</sub>-C<sub>8</sub>-dialkylaminocarbonyl and C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl,

R<sup>5</sup> and R<sup>8</sup> in each case independently of one another represent hydrogen, halogen or represent in each case optionally substituted C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, R<sup>12</sup>, G, J, -OJ, -OG, -S(O)<sub>p</sub>-J, -S(O)<sub>p</sub>-G, -S(O)<sub>p</sub>-phenyl, where the substituents independently of one another may be selected from one to three radicals W or from the group consisting of R<sup>12</sup>, C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy and C<sub>1</sub>-C<sub>4</sub>-alkylthio, where each substituent may be substituted by one or more substituents independently of one

- another selected from the group consisting of G, J, R<sup>6</sup>, halogen, cyano, nitro, amino, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl, phenyl and phenoxy, where each phenyl or phenoxy ring may optionally be substituted and where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R<sup>12</sup>,
- G in each case independently of one another represent a 5- or 6-membered non-aromatic carbocyclic or heterocyclic ring which may optionally contain one or two ring members from the group consisting of C(=O), SO and S(=O)<sub>2</sub> and which may optionally be substituted by one to four substituents independently of one another selected from the group consisting of C<sub>1</sub>-C<sub>2</sub>-alkyl, halogen, cyano, nitro and C<sub>1</sub>-C<sub>2</sub>-alkoxy, or independently of one another represent C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, (cyano)-C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, (C<sub>1</sub>-C<sub>4</sub>-alkyl)-C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, (C<sub>3</sub>-C<sub>6</sub>-cycloalkyl)-C<sub>1</sub>-C<sub>4</sub>-alkyl, where each cycloalkyl, (alkyl)cycloalkyl and (cycloalkyl)alkyl may optionally be substituted by one or more halogen atoms,
- J in each case independently of one another represent an optionally substituted 5- or 6-membered heteroaromatic ring, where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R<sup>12</sup>,
- R<sup>6</sup> independently of one another represent -C(=E<sup>1</sup>)R<sup>19</sup>, -LC(=E<sup>1</sup>)R<sup>19</sup>, -C(=E<sup>1</sup>)LR<sup>19</sup>, -LC(=E<sup>1</sup>)LR<sup>19</sup>, -OP(=Q)(OR<sup>19</sup>)<sub>2</sub>, -SO<sub>2</sub>LR<sup>18</sup> or -LSO<sub>2</sub>LR<sup>19</sup>, where each E<sup>1</sup> independently of one another represents O, S, N-R<sup>15</sup>, N-OR<sup>15</sup>, N-N(R<sup>15</sup>)<sub>2</sub>, N-S=O, N-CN or N-NO<sub>2</sub>,
- R<sup>7</sup> represents hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, halogen, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl,

- R<sup>9</sup> represents C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl or halogen,
- R<sup>10</sup> represents hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, halogen, cyano or C<sub>1</sub>-C<sub>4</sub>-haloalkoxy,
- R<sup>11</sup> in each case independently of one another represents in each case optionally mono- to trisubstituted C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfenyl, C<sub>1</sub>-C<sub>6</sub>-haloalkylthio, C<sub>1</sub>-C<sub>6</sub>-haloalkylsulfenyl, phenylthio or phenylsulfenyl, where the substituents independently of one another may be selected from the list W, -S(O)<sub>n</sub>N(R<sup>16</sup>)<sub>2</sub>, -C(=O)R<sup>13</sup>, -L(C=O)R<sup>14</sup>, -S(C=O)LR<sup>14</sup>, -C(=O)LR<sup>13</sup>, -S(O)<sub>n</sub>NR<sup>13</sup>C(=O)R<sup>13</sup>, -S(O)<sub>n</sub>NR<sup>13</sup>C(=O)LR<sup>14</sup> or -S(O)<sub>n</sub>NR<sup>13</sup>S(O)<sub>2</sub>LR<sup>14</sup>,
- L in each case independently of one another represents O, NR<sup>18</sup> or S,
- R<sup>12</sup> in each case independently of one another represents -B(OR<sup>17</sup>)<sub>2</sub>, amino, SH, thiocyanato, C<sub>3</sub>-C<sub>8</sub>-trialkylsilyloxy, C<sub>1</sub>-C<sub>4</sub>-alkyl disulfide, -SF<sub>5</sub>, -C(=E<sup>1</sup>)R<sup>19</sup>, -LC(=E<sup>1</sup>)R<sup>19</sup>, -C(=E<sup>1</sup>)LR<sup>19</sup>, -LC(=E<sup>1</sup>)LR<sup>19</sup>, -OP(=Q)(OR<sup>19</sup>)<sub>2</sub>, -SO<sub>2</sub>LR<sup>19</sup> or -LSO<sub>2</sub>LR<sup>19</sup>,
- Q represents O or S,
- R<sup>13</sup> in each case independently of one another represent hydrogen or represent in each case optionally mono- or polysubstituted C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, where the substituents independently of one another may be selected from the group consisting of R<sup>6</sup>, halogen, cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino or (C<sub>1</sub>-C<sub>4</sub>-alkyl)-C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino,
- R<sup>14</sup> in each case independently of one another represent in each case optionally mono- or polysubstituted C<sub>1</sub>-C<sub>20</sub>-alkyl, C<sub>2</sub>-C<sub>20</sub>-alkenyl, C<sub>2</sub>-C<sub>20</sub>-alkynyl or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, where the substituents independently of one another may be selected from the group consisting of R<sup>6</sup>, halogen, cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino and (C<sub>1</sub>-C<sub>4</sub>-alkyl)-C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino or represent optionally substituted phenyl, where


- the substituents independently of one another may be selected from one to three radicals W or one or more radicals R<sup>12</sup>,
- R<sup>15</sup> in each case independently of one another represent hydrogen or represent in each case optionally mono- or polysubstituted C<sub>1</sub>-C<sub>6</sub>-haloalkyl or C<sub>1</sub>-C<sub>6</sub>-alkyl, where the substituents independently of one another may be selected from the group consisting of cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl and optionally substituted phenyl, where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R<sup>12</sup>, or N(R<sup>15</sup>)<sub>2</sub> represents a cycle which forms the ring M,
- R<sup>16</sup> represents C<sub>1</sub>-C<sub>12</sub>-alkyl or C<sub>1</sub>-C<sub>12</sub>-haloalkyl, or N(R<sup>16</sup>)<sub>2</sub> represents a cycle which forms the ring M,
- R<sup>17</sup> in each case independently of one another represent hydrogen or C<sub>1</sub>-C<sub>4</sub>-alkyl, or B(OR<sup>17</sup>)<sub>2</sub> represents a ring in which the two oxygen atoms are attached via a chain having two to three carbon atoms which are optionally substituted by one or two substituents independently of one another selected from the group consisting of methyl and C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl,
- R<sup>18</sup> in each case independently of one another represent hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl or C<sub>1</sub>-C<sub>6</sub>-haloalkyl, or N(R<sup>13</sup>)(R<sup>18</sup>) represents a cycle which forms the ring M,
- R<sup>19</sup> in each case independently of one another represent hydrogen or represent in each case mono- or polysubstituted C<sub>1</sub>-C<sub>6</sub>-alkyl, where the substituents independently of one another may be selected from the group consisting of cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, CO<sub>2</sub>H, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl and optionally substituted phenyl, where the substituents

- independently of one another may be selected from one to three radicals W, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl or phenyl or pyridyl, each of which is optionally mono- to trisubstituted by W,
- M in each case represents an optionally mono- to tetrasubstituted ring which, in addition to the nitrogen atom attached to the substituent pair R<sup>13</sup> and R<sup>18</sup>, (R<sup>15</sup>)<sub>2</sub> or (R<sup>16</sup>)<sub>2</sub>, contains two to six carbon atoms and optionally additionally a further nitrogen, sulfur or oxygen atom, where the substituents independently of one another may be selected from the group consisting of C<sub>1</sub>-C<sub>2</sub>-alkyl, halogen, cyano, nitro and C<sub>1</sub>-C<sub>2</sub>-alkoxy,
- W in each case independently of one another represent C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>2</sub>-C<sub>4</sub>-haloalkenyl, C<sub>2</sub>-C<sub>4</sub>-haloalkynyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, halogen, cyano, nitro, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, (C<sub>1</sub>-C<sub>4</sub>-alkyl)-C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, C<sub>2</sub>-C<sub>4</sub>-alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, CO<sub>2</sub>H, C<sub>2</sub>-C<sub>6</sub>-alkylaminocarbonyl, C<sub>3</sub>-C<sub>8</sub>-dialkylaminocarbonyl or C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl,
- n in each case independently of one another represent 0 or 1,
- p in each case independently of one another represent 0, 1 or 2,
- where, if (a) R<sup>5</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkynyl, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio or halogen and (b) R<sup>8</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkynyl, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, halogen, C<sub>2</sub>-C<sub>4</sub>-alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkylaminocarbonyl or C<sub>3</sub>-C<sub>8</sub>-dialkylaminocarbonyl, (c) at least one substituent selected from the group consisting of R<sup>6</sup>, R<sup>11</sup> and R<sup>12</sup> is present and (d) if R<sup>12</sup> is not present, at least one of the radicals R<sup>6</sup> and R<sup>11</sup> is different from C<sub>2</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkylaminocarbonyl and C<sub>3</sub>-C<sub>8</sub>-dialkylaminocarbonyl, and where the compound of the general formula (I) may also be an N-oxide or salt,

and at least one insecticidally active compound of groups 2 and 3 below selected from


A) (group 2),

(2-1) azinphos-methyl




and/or

(2-2) chlorpyrifos




and/or

(2-3) diazinon




and/or

(2-4) dimethoate




and/or

(2-5) disulfoton




and/or

(2-6) ethion




and/or

(2-7) fenitrothion




~~ad/or~~ and/or

(2-8) fenthion




and/or

(2-9) isoxathion




and/or

(2-10) malathion



and/or


(2-11) methidathion



and/or


(2-12) oxydemeton-methyl






and/or

(2-13) parathion




and/or

(2-14) parathion-methyl




and/or

(2-15) phenthoate




and/or

(2-16) phorate




and/or

(2-17) phosalone




and/or

(2-18) phosmet




and/or

(2-19) phoxim




and/or

(2-20) pirimiphos-methyl




and/or

(2-21) profenophos




and/or

(2-22) prothiophos




and/or

(2-23) tebupirimphos




and/or

(2-24) triazophos




and/or

(2-25) chlorfenvinphos




and/or

(2-26) dichlorophos




and/or

(2-27) dicrotophos




and/or

(2-28) mevinphos




and/or

(2-29) monocrotophos




and/or

(2-30) phosphamidon




and/or

(2-31) acephate




and/or

(2-32) methamidophos



and/or


(2-33) trichlorfon



and/or


B) (group 3),

(3-1) carbaryl




and/or

(3-2) fenoxycarb



and/or

(3-3) formetanate




and/or

(3-4) formetanate hydrochloride


and/or

(3-5) methiocarb




and/or

(3-6) methomyl




and/or

(3-7) oxamyl




and/or

(3-8) pirimicarb (= Pirimor)




and/or

(3-9) propoxur




and/or

(3-10) thiodicarb



2. (Currently amended) The composition as claimed in claim 1 comprising at least one compound of the formula (I-1) ~~in which~~:



in which

- $R^2$  represents hydrogen or  $C_1$ - $C_6$ -alkyl,  
 $R^3$  represents  $C_1$ - $C_6$ -alkyl which is optionally substituted by one  $R^6$ ,  
 $R^4$  represents  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_2$ -haloalkyl,  $C_1$ - $C_2$ -haloalkoxy or halogen,  
 $R^5$  represents hydrogen,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_2$ -haloalkyl,  $C_1$ - $C_2$ -haloalkoxy or halogen,  
 $R^6$  represents  $-C(=E^2)R^{19}$ ,  $-LC(=E^2)R^{19}$ ,  $-C(=E^2)LR^{19}$  or  $-LC(=E^2)LR^{19}$ , where each  $E^2$  independently of one another represents O, S,  $N-R^{15}$ ,  $N-OR^{15}$ ,  $N-N(R^{15})_2$ , and each L independently of one another represents O or  $NR^{18}$ ,  
 $R^7$  represents  $C_1$ - $C_4$ -haloalkyl or halogen,  
 $R^9$  represents  $C_1$ - $C_2$ -haloalkyl,  $C_1$ - $C_2$ -haloalkoxy,  $S(O)_p$ - $C_1$ - $C_2$ -haloalkyl or halogen,

R<sup>15</sup> in each case independently of one another represent hydrogen or represent in each case optionally substituted C<sub>1</sub>-C<sub>6</sub>-haloalkyl or C<sub>1</sub>-C<sub>6</sub>-alkyl, where the substituents independently of one another may be selected from the group consisting of cyano, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl and C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl,

R<sup>18</sup> in each case represents hydrogen or C<sub>1</sub>-C<sub>4</sub>-alkyl,

R<sup>19</sup> in each case independently of one another represent hydrogen or C<sub>1</sub>-C<sub>6</sub>-alkyl,

and

p independently of one another represent 0, 1, 2.

3. (Previously presented) The composition as claimed in claim 1 or 2 comprising at least one compound selected from the group consisting of
  - (2-2) chlorpyrifos,
  - (2-31) acephate,
  - (2-32) methamidophos,
  - (3-1) carbaryl,
  - (3-5) methiocarb, and
  - (3-10) thiodicarb.
4. (Previously presented) The composition as claimed in claim 1, comprising at least one compound of formula (I) and at least one compound from group 2 and/or group 3 in a ratio of 50:1 to 1:50.
5. (Cancelled).
6. (Previously presented) A process for preparing pesticides, comprising contacting a synergistically effective mixture as defined in claim 1 with extenders and/or surfactants.

7. (Previously presented) A method for controlling animal pests, comprising allowing a synergistically effective mixture as defined in claim 1 to act on animal pests and/or their habitat.